

Cecil, F.E. and G.M. Hale. *Measurement of D-D and D-Li⁶ Nuclear Reactions at Very Low Energies*. in *Second Annual Conference on Cold Fusion, "The Science of Cold Fusion"*. 1991. Como, Italy: Societa Italiana di Fisica, Bologna, Italy.

MEASUREMENT OF D-D AND D-⁶LI NUCLEAR REACTIONS AT VERY LOW ENERGIES

F.E. Cecil^a and G.M. Hale^b

^aColorado School of Mines, Golden CO 80401, USA

^bLos Alamos National Laboratory, Los Alamos NM 87545, USA

ABSTRACT

The nuclear reactions of very low energy deuterons (down to center-of-mass energies of 2 keV) with deuterons and ⁶Li have been measured. The measured D-D reactions are in good with agreement recent R-matrix calculations. The reaction ratios D(d,p)T/D(d,n)³He and ⁶Li(d,p)⁷Li/⁶Li(d,α)⁴He in particular were examined for possible evidence of an Oppenheimer-Phillips type enhancement. No significant enhancement was found in either ratio or in the absolute yields of the reactions. The radiative capture reactions D(d,γ)⁴He and ⁶Li(d,γ)⁸Be were likewise measured. The branching ratios of these radiative capture reactions to the nucleonic branches of the reactions appear roughly independent of energy. The role of these reactions in the production of heat in cold-fusion experiments is evaluated.

INTRODUCTION

The nuclear reactions between two deuterons and perhaps between a deuteron and a ⁶Li nucleus are generally accepted as playing crucial roles in recently observed nuclear processes and significant heat production in condensed matter deuterium-metal systems. Independent measurements of the cross sections for these nuclear reactions at low energies will allow these assumed roles to be addressed. The ratio of the reactions D(d,n)³He and D(d,p)T or the ratio of the reactions D(d,γ)⁴He and D(d,p)T at very low energies will, for example, determine whether significant heat production is possible from D-D nuclear reactions in the absence of enormous quantities of escaping and potentially hazardous radiation. Similarly, the ratio of the reactions ⁶Li(d,p)⁷Li and ⁶Li(d,α)⁴He and the ratio of the reactions ⁶Li(d,γ)⁸Be and ⁶Li(d,α)⁴He will allow the analogous determination for the D-⁶Li reaction to be addressed. In this work, we report on our recent measurements of these reactions. While some of these reactions have been studied at relatively low energies¹⁻⁵, the present work extends our knowledge of these reactions to significantly lower energies. On the other hand, our measurement of the reaction ⁶Li(d,γ)⁸Be is the first reported observation of this reaction.

EXPERIMENTAL RESULTS

The D-D and D-⁶Li reactions were studied using magnetically analyzed deuteron beams from the Colorado School of Mines low energy charged particle accelerator⁶. The targets consisted of pressed sheets of CD₂ and rolled foils of metallic Li, isotopically enriched to 94% ⁶Li. The charged reaction products were detected with silicon surface barrier detectors placed at 150° from the beam direction and protected from the Rutherford backscattered beam deuterons by a thin Al foil. The gamma rays were detected with a NaI(Tl) scintillation spectrometer surrounded by an active Compton scattered gamma ray and cosmic ray shield. This gamma ray detector system has been described elsewhere⁷. The techniques used in the gamma ray to charged particle branching ratio measurements have likewise been described in some of our earlier studies⁸.

Charged particle spectra measured during the bombardment of the CD₂ and ⁶Li targets are shown in Figure 1.




Figure 1. (a) Charged particle spectrum for D-D reaction at $E_{\text{lab}} = 10$ keV. (b) Charged particle spectrum for $D\text{-}^6\text{Li}$ reaction at $E_{\text{lab}} = 150$ keV.

Since the angular distributions of the outgoing reaction products for the D-D and $D\text{-}^6\text{Li}$ reactions have been shown to be nearly isotropic at low energies¹⁻³, the reaction ratios $D(d,p)T/D(d,n)^3\text{He}$ and $^6\text{Li}(d,p)^7\text{Li}/^6\text{Li}(d,\alpha)^4\text{He}$ are determined directly from the ratio of the yield of the peaks labeled “ ^3He ions” and “tritons” for the D-D reaction and “p0” and “ α ” for the $D\text{-}^6\text{Li}$ reaction. The yield ratios so determined as a function of energy are given in Figure 2. The yield ratios for the D-D reaction presented in Figure 2 are compared to a recent R-matrix⁹ calculation. Our measured yield ratios are consistent with the calculated ratio, indicating no enhancement of the (d,p) reaction as qualitatively suggested by Oppenheimer and Phillips¹⁰. While there have been no comparable calculations for the ratio of the reactions $^6\text{Li}(d,p)^7\text{Li}$ and $^6\text{Li}(d,\alpha)^4\text{He}$, Koonin¹¹ has calculated the (d,p)/(d,n) ratio for the $D\text{-}^6\text{Li}$ reactions at low energies and has predicted no significant enhancement of the (d,p) reaction. Our results are consistent with these predictions since we would expect the (d, α) reaction to be unaffected by any Oppenheimer-Phillips type processes.

These spectra also allowed a determination of the absolute thick target yield of the reactions. The measured yields are shown in Figure 3 and are compared, respectively, to yields based upon the R-matrix calculation noted above or calculated assuming a slowly varying astrophysical S-factor. There is good agreement between measured and calculated yields again indicating no anomalous behavior at very low energies.




Figure 2. (a) Ratio of (d,p) to (d,n) branches for D-D reactions, (b) Ratio of (d,p) to (d, α) branches for $D\text{-}^6\text{Li}$ reactions.

Gamma ray spectra were measured during the deuteron bombardment of the CD_2 and ^6Li targets between c.m. energies of 20 and 100 keV. The yield of the 23.8 MeV and 22.6 MeV gamma rays and the concurrent yield of the charged particles (see Figs 1) will determine the gamma ray to charged particle branching ratios for the D-D and $D\text{-}^6\text{Li}$ reactions respectively after the yield ratios are corrected for the relative detector efficiencies. The branching ratios so determined are plotted in Figure 4. The D-D branching ratio is consistent at the higher energies with our earlier measurements. The fact that the measurements of the D-D gamma ray to charged particle branching ratio at c.m. energies of 20 and 40 keV are, to within errors, equal in value, suggest that the branching ratio is independent of energy. The energy dependence for the branching ratio for the $D\text{-}^6\text{Li}$ reaction is somewhat inconclusive although energy independence is certainly not ruled out by virtue of the relatively large error bar on the lower energy data point for this reaction.




Figure 3. (a) Measured and calculated yields for D(d,p)T reaction, (b) Measured and calculated yields for ${}^6\text{Li}(d,\alpha)^4\text{He}$ reaction.




Figure 4. Gamma ray to charged particle branching ratios for the D-D and D- ${}^6\text{Li}$ reactions. The open squares are from Ref. 4.

CONCLUSIONS

The results of our measurements have significant implications for any effort to associate the D-D or D- ${}^6\text{Li}$ nuclear reactions with reports of heat production from cold fusion experiments. Specifically, if the particle-particle and gamma ray-particle branching ratios which we have measured at very low laboratory energies are characteristic of the branching ratios occurring in condensed matter fusion in deuterium-metal systems, then even low levels of heat production arising from these nuclear reactions will necessarily be associated with enormous quantities of escaping radiation. For example, based on the branching ratios given in Figures 2 and 4, if 1 Watt of power were produced by the D-D reaction, then there would be an accompanying production of about 10^{12} 2.5 MeV neutrons per second and about 10^8 24 MeV gamma rays per second. Comparable yields of neutrons and gamma rays will be associated with similar levels of power production by the D- ${}^6\text{Li}$ reaction. If, therefore, the reported production of heat from condensed matter fusion in deuterium-metal systems is to be attributed to D-D or D- ${}^6\text{Li}$ reactions, then the particle-particle and gamma ray-particle branching ratios at unmeasurably low energies must vary drastically from those measured at the low energies reported in this work.

This work is supported by the U.S. Department of Energy.

References

1. R. Brown and N. Jarmie, Phys. Rev. C41 (1990) 1391.
2. A. Krauss et al, Nucl. Phys. A465 (1987) 150.
3. A.J. Elwyn et al., Phys. Rev. C16 (1977) 1744.
4. F.J. Wilkinson and F.E. Cecil, Phys. Rev. C31 (1985) 2036.
5. C.A. Barnes et al., Phys. Lett. 197B (1987) 1922.
6. F.E. Cecil et al., NIM B40/41 (1989) 934.
7. F.E. Cecil et al., NIM A234 (1985) 479.
8. F.E. Cecil and F.J. Wilkinson, Phys. Rev. Letts. 53 (1984) 767.
9. G.M. Hale and D.C. Dodder, Proc. Int. Conf. on Nucl. Cross Sections for Technology. Knoxville (1979) J.C. Fowler, C.H. Johnson and C.D. Bowman editors. NBS Special Publication 594, p. 650.
10. J.R. Oppenheimer and M. Phillips, Phys. Rev. 48 (1935) 500.
11. S.E. Koonin and M. Mukerjee, Phys. Rev. C42 (1990) 1639.